



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Let $\dot{\theta}$ = the angular motion of the instantaneous axis on the plane; this is the angular motion about the line (n) which is normal to the inclined plane and passes through the vertex of the cone. A plane perpendicular to the inclined plane includes this normal, the instantaneous axis (a_i), the axis of the cone, and a perpendicular (p) to the axis of the cone. The angle between n and p is α and between a_i and p is $\pi/2 + \alpha$. The components of $\dot{\theta}$ and ω about p are $\dot{\theta} \cos \alpha$ and $\omega \cos (\pi/2 + \alpha) = -\omega \sin \alpha$, but these denote the same motion; then

$$\dot{\theta} \cos \alpha = -\omega \sin \alpha \quad \text{or} \quad \omega = -\dot{\theta} \cot \alpha. \quad (6)$$

Now the moment of the external forces about the instantaneous axis

$$= L \cos \lambda + M \cos \mu + N \cos \nu; \quad (7)$$

also

$$= \frac{3}{4} mgh \sin \alpha \sin \beta \sin \theta \quad (8)$$

β being the inclination of the plane.

Substituting (4) in (3) and differentiating, we have

$$\dot{\omega}_1 = \sin \alpha (\cos \varphi \cdot \dot{\varphi}^2 + \sin \varphi \cdot \ddot{\varphi}), \quad \dot{\omega}_2 = \sin \alpha (-\sin \varphi \cdot \dot{\varphi}^2 + \cos \varphi \cdot \ddot{\varphi}), \quad \text{and} \quad \dot{\omega}_3 = \cos \alpha \cdot \ddot{\varphi}, \quad (9)$$

Now put the values given by (3), (5), (9) in (1), and we have the values of L , M , N ; and using these, with (2), (7) and (8), we have the single equation of motion

$$\frac{3m}{20} (6 + \tan^2 \alpha) h^2 \sin^2 \alpha \cdot \ddot{\varphi} = \frac{3m}{4} gh \sin \alpha \sin \beta \sin \theta. \quad (10)$$

Now from (4) and (6),

$$\ddot{\theta} = -\frac{g}{h} \frac{5 \sin \beta \cos \alpha}{1 + 5 \cos^2 \alpha} \sin \theta. \quad (11)$$

If θ be so small that θ may displace $\sin \theta$, we have the required time of oscillation

$$T = 2\pi \sqrt{\frac{1 + 5 \cos^2 \alpha}{5 \sin \beta \cos \alpha} \cdot \frac{h}{g}}.$$

Also solved by F. L. WILMER, and discussed by C. H. ECKART.

2820 [1920, 134]. Proposed by C. B. HALDEMAN, Ross, Ohio.

Given one angle and the radii of the inscribed and circumscribed circles, to construct the triangle geometrically.

SOLUTION BY A. V. RICHARDSON, Bishop's College, Lennoxville, Quebec.

With the usual notation, A , R , r are given. Draw a circle of radius R , and let KB be any diameter: (The reader is requested to draw the figure). Make the angle, $BKC = A$. Draw XY parallel to BC and at a distance r from it on the same side as K . Bisect the arc BC at V , and let the circle, center V , radius VC , cut XY at I and I' .

These points will be the incenters for the two (symmetrical) solutions. Let VI meet the circle again in A . Then $\triangle ABC$ is the required triangle. To show this, it is only necessary to prove that the straight line CI bisects the angle C . Since $VI = VC$, $\angle VIC = \angle VCI$. But $\angle VIC = (A/2) + \angle ICA$ and $\angle VCI = (A/2) + \angle BCI$. Hence $\angle ICA = \angle BCI$ and I is the intersection of two bisectors of the angles of the triangle ABC .

Also solved by L. C. MATHEWSON, H. L. OLSON, ARTHUR PELLETIER, J. B. REYNOLDS, JOSEPH ROSENBAUM, C. N. SCHMALL, and the Proposer.

2829 [1920, 226]. Proposed by E. S. PALMER, New Haven, Conn.

Given a set of arbitrary pairs of positive integers (a_p, b_p), ($p = 1, 2, \dots, n$): (a) Is it always possible to find a set of positive integers k_p , ($p = 1, 2, \dots, n$) such that

$$k_p a_p + k_p b_p > \sum_{r=1}^{r=n} k_r a_r, \quad (p = 1, 2, 3, \dots, n).$$

(b) If or when possible, show how to find k_p .